

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
7 October 2004 (07.10.2004)

PCT

(10) International Publication Number
WO 2004/085477 A1

(51) International Patent Classification⁷: **C07K 16/24**,
A61K 39/395, 39/385, 39/00, A61P 37/02, 11/00, 37/08

(21) International Application Number:
PCT/EP2004/050375

(22) International Filing Date: 26 March 2004 (26.03.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
03447070.8 28 March 2003 (28.03.2003) EP
60/461,687 9 April 2003 (09.04.2003) US

(71) Applicant (for all designated States except US): INNO-
GENETICS N.V. [BE/BE]; Technologiepark 6, B-9052
Ghent (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DEPRAETERE**,
Stany [BE/BE]; Gentstraat 224, B-9700 Oudenaarde
(BE). **LORRÉ**, Katrien [BE/BE]; Lievekaai 12, B-9000
Ghent (BE). **BUYSE**, Marie-Ange [BE/BE]; Burg. E.
Ronsestraat 23, B-9820 Merelbeke (BE).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,
GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments
- with (an) indication(s) in relation to deposited biological
material furnished under Rule 13bis separately from the
description

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: TREATMENT OF TYPE 1 IMMUNE RESPONSE-MEDIATED INFLAMMATORY LUNG DISEASE BY MODULA-
TION OF IFN-GAMMA ACTIVITY

(57) Abstract: The present invention relates to preventing, or treating and/or reducing the severity or progression of Type 1 immune response-mediated inflammatory lung disease. More particularly, the present invention provides a method for preventing or treating chronic obstructive pulmonary disease (COPD), severe asthma, sarcoidosis, berylliosis or cystic fibrosis by neutralizing or reducing IFN γ bioactivity which can be achieved either by in vivo administration of IFN γ neutralizing molecules or by in vivo immunization with pharmaceutical compositions comprising immunogenic IFN γ proteins or IFN γ -derived (poly)peptides or their corresponding nucleic acid sequences.



WO 2004/085477 A1

10/550175
JC17 Rec'd PCT/PTO 21 SEP 2005

1/PRTS

1

TREATMENT OF TYPE 1 IMMUNE RESPONSE-MEDIATED
INFLAMMATORY LUNG DISEASE BY
MODULATION OF IFN-GAMMA ACTIVITY

FIELD OF THE INVENTION

The present invention relates to preventing, or treating and/or reducing the severity or progression of Type 1 immune response-mediated inflammatory lung disease. More particularly, the present invention provides a method for preventing or treating chronic obstructive pulmonary disease (COPD), severe asthma, sarcoidosis, berylliosis or cystic fibrosis by neutralizing or reducing IFN γ bioactivity which can be achieved either by in vivo administration of IFN γ neutralizing molecules or by in vivo immunization with pharmaceutical compositions comprising immunogenic IFN γ proteins or IFN γ -derived (poly)peptides or their corresponding nucleic acid sequences.

BACKGROUND ART

The present invention relates to preventing the onset of symptoms, treating and/or reducing the severity or progression of COPD and other Type 1 immune response-mediated (T1) inflammatory lung diseases such as, but not limited to, severe asthma, sarcoidosis, berylliosis, and cystic fibrosis. T1 inflammatory lung diseases are characterized by a Type 1 immune response mediated by T helper-1 cells (CD4+) and T cytotoxic-1 cells (CD8+) and by increased production of interferon gamma (IFN γ), tumor necrosis factor (TNF), and interleukin-2 (IL-2). T1 cytokines evoke cell-mediated immunity characterized by prominent lung tissue infiltration of macrophages, neutrophils, and T-cells.

Severe asthma

Asthma is a disease of the respiratory system characterized by hyper-responsiveness to bronchoconstricting stimuli, inflammation, and changes in respiratory epithelium. Asthmatic patients in whom the disease process is either refractory to therapy or

requires persistent use of high dose systemic anti-inflammatory corticosteroids in order to maintain reasonable control of symptoms are referred to in literature as patients suffering from severe or irreversible or refractory asthma (Kaplan N.M. et al., 2000). Patients with severe persistent asthma have continual symptoms, frequent exacerbations, frequent nighttime symptoms and evidence of severe obstructive lung disease on pulmonary function testing (Forced Expiratory Volume in one second: FEV1<60%). The narrowing of airways causes ventilation perfusion imbalance, lung hyperventilation, and increased work of breathing that may lead to ventilatory muscle fatigue and life-threatening respiratory failure (Papiris S. et al., 2002). WO 01/34180 by Block Lutz-Henning describes the use of IFN- γ for the treatment of severe asthma bronchiale. New effective treatments are needed for this subpopulation of asthmatic patients (Stirling R.G. and Chung K.F., 2001).

Increased numbers of neutrophils are observed in the lung mucosa and the bronchoalveolar fluid of severe asthmatics (Wenzel et al., 1997). Eicosanoid mediators such as thromboxane and leukotriene B4 are also high in the lung tissue of these patients. No difference in eosinophil concentration is observed in BAL fluid derived from healthy controls or from severe asthmatics.

Severe asthma is differentiated from mild/moderate asthma by distinct inflammatory processes involving varied cytokine expression profiles and/or effector cells. In contrast to severe asthmatics, mild or moderate asthmatics show enhanced concentrations of eosinophils in their lungs and a Type 2 inflammation characterized by predominant infiltration of T helper-2 T lymphocytes that produce IL-4, IL-5, IL-9 and IL-13. Neutrophils are absent. Symptoms in mild to moderate asthmatics are well controlled by treatment with β 2-agonists and corticosteroids.

Sarcoidosis

Sarcoidosis and berylliosis are interstitial lung diseases. The interstitium (the space between the tissues) of the lungs includes portions of the connective tissue of the blood vessels and air sacs. Interstitial lung diseases begin with inflammation of the lung cells. The lungs stiffen as a result of inflammation of the air sacs (alveolitis) and scarring (fibrosis) (The Lungs in Health and Disease, National Heart, Lung and Blood Institute; NIH Publication No. 97-3279, August 1997).

Sarcoidosis is a systemic granulomatous disease of unknown aetiology and world-wide distribution. It most commonly affects young adults and presents as with bilateral hilar lymphadenopathy, pulmonary infiltration, reticuloendothelial involvement, eye and skin lesions. Shortness of breath (dyspnea) and a cough that will not go away can be among the first symptoms of sarcoidosis (Sarcoidosis, National Heart, Lung, and Blood Institute; NIH publication No. 95-3093, reprinted July 1995). The hallmark of pulmonary sarcoidosis is a mononuclear alveolitis which is characterized by activated CD4+ lymphocytes, monocytes/macrophages, and non-caseating granulomas. An imbalance in the expression of T1 and T2 cytokines by alveolar cells is thought to play an important role in the immunopathogenesis of sarcoidosis. It is well established that T1 cytokines are important mediators in pulmonary sarcoidosis and that there is a dependence of granulomatous inflammation on T1 cytokines. Alveolar cells spontaneously release the T1 cytokines IFN γ and IL-2, but not T2 cytokines. However, it is suggested that the cytokine patterns change during the course of the disease (Möllers M., et al., 2001). Shigehara K. et al., 2001, demonstrated that IL-12 and IL-18 were increased in BAL fluids of patients with sarcoidosis. IL-12 and IL-18 drive the immune response in to Type 1 direction. In recent years, new therapies have been studied for sarcoidosis. Drugs used to treat patients with sarcoidosis are corticosteroids, cytotoxic agents, immunomodulators (chloroquine and hydroxychloroquine) and the antileprosy drugs clofazimine and minocycline. A key cytokine in chronic sarcoidosis appears to be TNF. Drugs that inhibit its release or block its bioactivity such as pentoxifylline and thalidomide have been reported to be effective for treatment of sarcoidosis (Baughman R.P., 2002).

Berylliosis

Berylliosis or Chronic Beryllium Disease (CBD) is an environmental chronic inflammatory disorder of the lungs caused by inhalation of insoluble beryllium (Be) dust and characterized by the accumulation of CD4+ T cells and macrophages in the lower respiratory tract. According to the type and level of Be exposure, the reactions vary from acute tracheobronchitis, chemical pneumonitis, and metal fume fever to a chronic granulomatous lung disorder. Lung T cells respond to beryllium with the release of T1 cytokines such as IL-2, the migration inhibitor factor (MIF), IFN γ , and TNF- α . Furthermore, a positive association with a HLA class II allele, HLA-DP, has

been described (Saltini C. et al., 2001). Most patients treated with corticosteroids need to remain on therapy for life (Rossman M.D., 2001).

Cystic Fibrosis

Cystic fibrosis (CF) is an autosomal recessive disorder caused by nearly 1000 different mutations of the cystic fibrosis transmembrane conductance regulator (CFTR) gene. It is a multisystem disorder characterized by defective electrolyte transport in epithelial cells and abnormally viscous mucus secretions from glands and mucus epithelia. Symptoms are pancreatic insufficiency (PI) associated with neonatal meconium ileus and chronic obstructive lung disease superimposed with recurrent opportunistic infections that progressively destroy lung tissue. The inflammatory response is a primary cause of irreversible lung damage. Inflammation is present in CF patients and precedes the bacterial infections, as demonstrated by increased levels of neutrophils, TNF, and IL-8. Other complications include liver disease, chronic sinusitis, infertility in male patients and elevated sweat chloride concentrations. Despite advances in genomic technologies and drug discovery, drug therapy often improves disease symptoms but does not cure the disease. One of the main causes of this failure to cure CF may be attributable to genetic variability and to the scarce knowledge of CF biochemistry. The development of new treatments may be important for the life expectancy of patients. Current CF therapeutic strategies include lung transplantation, antimicrobial treatment, corticosteroids, non-steroidal anti-inflammatory drugs like ibuprofen, ion channel therapy, protein-assist therapy, and gene therapy (Sanguolo F. et al., 2002).

COPD

Chronic obstructive pulmonary disease (COPD) is currently the sixth leading cause of death and the 12th leading cause of morbidity world-wide. By the year 2020, COPD is expected to be the third leading cause of death and the fifth leading cause of disability.

COPD is a disease state characterized by chronic and slow progressive development of airflow limitation that is not fully reversible and is punctuated by episodic exacerbations due to viral or bacterial infections. The airflow limitation is associated with an abnormal inflammatory response of the lungs to noxious particles or gases (Executive Summary, Global Strategy for the diagnosis, management and prevention

of chronic obstructive pulmonary disease; NHLBI/WHO Workshop Report). COPD comprises chronic bronchitis, chronic obstructive bronchiolitis and emphysema (Leckie M.J. et al., 2000). Chronic bronchitis is defined as the occurrence of coughing and by production of sputum on most days for at least 3 months over 2 consecutive years. Emphysema is characterized by destructive enlargement of airspaces with loss of normal architecture and lung elasticity.

Cigarette smoking is the dominant factor for the development and progression of COPD. However, only 15% of smokers develop COPD and >15% of COPD-related mortality occurs in people who have never smoked, suggesting that other factors are important. Genetic factors such as α_1 -antitrypsin deficiency, resulting in enhanced neutrophil elastase activity, account for 2% of emphysema patients and polymorphism of the TNF- α gene, leading to enhanced TNF production, may also play an important role. The role of infections in both the development and progression of COPD is getting increased attention, including adenoviral and rhinoviral infections in patients with emphysema. Occupational and environmental exposures to various pollutants are also considered to be important factors in the development of COPD. (Mannino D.M. et al., 2002; Barnes P.J., 2000).

A diagnosis of COPD should be considered in any patient who has symptoms of cough, sputum production, or dyspnea, and/or a history of exposure to risk factors for the disease. Exhaled gases (nitric oxide and carbon monoxide) and inflammatory markers in exhaled breath condensate can be used as non-invasive markers of COPD (Leckie M.J. et al., 2000). The diagnosis is confirmed by spirometry. The presence of a post-bronchodilator $FEV_1 < 80\%$ of the predicted value in combination with an $FEV_1/FVC < 70\%$ confirms the presence of airflow limitation that is not fully reversible (FEV_1 = forced expiratory volume in one second; FVC = forced vital capacity).

A simple classification of disease severity into four stages is recommended. All FEV_1 values refer to post-bronchodilator FEV_1 .

- *Stage 0: At Risk:* characterized by chronic cough and sputum production. Lung function, as measured by spirometry, is still normal.

- *Stage I: Mild COPD*: characterized by mild airflow limitation ($FEV_1/FVC < 70\%$ but FEV_1 80% predicted) and usually, but not always, by chronic cough and sputum production.
- *Stage II: Moderate COPD*: characterized by worsening airflow limitation ($30\% \leq FEV_1 < 80\%$ predicted) and usually the progression of symptoms, with shortness of breath typically developing on exertion. This is the stage at which patients typically seek medical attention because of dyspnea or an exacerbation of their disease. The division into stages IIA and IIB is based on the fact that exacerbations are especially seen in patients with an FEV_1 below 50% predicted. The presence of repeated exacerbations has an impact on the quality of life of patients and requires appropriate management.
- *Stage III: Severe COPD*: characterized by severe airflow limitation ($FEV_1 < 30\%$ predicted) or the presence of respiratory failure or clinical signs of right heart failure. Patients may have severe (Stage III) COPD even if the FEV_1 is $> 30\%$ predicted, whenever these complications are present. At this stage, quality of life is appreciably impaired and exacerbations may be life-threatening.

COPD is generally regarded as a separate condition from asthma (reversible airflow limitation) in terms of inflammatory processes, underlying pathology, and responses to treatment. Airway inflammation in asthma is characterized by infiltration of eosinophils and T helper-2 lymphocytes. Macrophages are less frequent and CD8 T cells are usually absent. The T2 cytokines predominate: IL-4 and IL-13 play an important role in IgE production, whereas IL-5 is critical for eosinophil growth and differentiation (Barnes P.J., 2000).

Airway inflammation in COPD is characterized by the presence of neutrophils, macrophages and CD8 T cells. Histopathological studies show that most inflammation in COPD occurs in the peripheral airways (bronchioles) and lung parenchyma. The bronchioles are obstructed by fibrosis and there is destruction of lung parenchyma. Bronchial biopsy results show similar findings. There is also a marked increase in macrophages and neutrophils observable in bronchoalveolar lavage fluid and induced sputum.

Emphysema is caused by an imbalance of proteases and protease inhibitors. The concentration of inflammatory mediators such as leukotriene B_4 , $TNF-\alpha$ and IL-8 are

increased in sputum of patients with COPD (Barnes P.J., 2000). Other inflammatory cytokines associated with COPD are TGF- β , IL-1, IL-6, IL-11 and IL-18. Also a role for IFN γ in COPD has been mentioned in the literature, however with some differing data. Wang Z. et al. (2000) describe a transgenic mouse model, with IFN γ inducibly targeted to the adult murine lung, showing emphysema, macrophage and neutrophil infiltration and inversed protease/anti-protease ratio. Majori M. et al. (1999) showed an increase in the percentage of IFN γ -producing cells among peripheral blood T helper cells from patients with COPD. Analysis of cytokine levels in BAL (bronchoalveolar lavage) discloses a prevalent T1 cytokine pattern in COPD, however this difference was not significant when COPD patients were compared with the other two groups (asthmatics and non-smokers) (Balbi B. et al., ATS 2002). Lethbridge M.W.G. et al. (2002) even showed a disproportionately low number of IFN γ -expressing airway lymphocytes in COPD smokers as compared to healthy smokers and healthy ex-smokers. The association of chronic inflammation with the pathophysiology of COPD makes IL-1, IL-18, and TNF- α targets for therapeutic intervention (de Boer W.L., 2002). Macrophages predominate and appear to play a central role as they have the capacity to produce all the pathologic changes of COPD (Hautamaki, 1997). Macrophages release LTB4 and IL-8, which are potent neutrophil chemo-attractants, and multiple proteases responsible for the continued proteolytic activity in the lungs of patients with emphysema.

At the present time, there are no known drugs that slow the relentless progression of COPD and there is a pressing need to develop new drugs to control the inflammatory and destructive processes that underlie the disease. Smoking cessation is the only measure that will slow the progression of COPD. However, even if the patient stops smoking, the damage already caused will continue to cause symptoms (Barnes P.J., 2001). Currently available drugs that provide symptomatic relief in COPD are:

- Bronchodilators, which are the mainstay of current drug therapy for COPD (Leckie M.J. et al., 2000).
- Inhaled corticosteroids are also widely prescribed for COPD but have a risk of systemic side effects (Barnes P.J., 2000). The usefulness of inhaled corticosteroids in COPD remains controversial (Leckie M.J. et al., 2000).
- Theophylline

- Although antibiotics are still widely used for exacerbations of COPD, it is increasingly recognized that exacerbations may be due to viral infections of the upper respiratory tract or may be noninfective, so that antibiotic treatment is not always warranted (Barnes P.J., 2002).

Nonpharmacologic treatments include oxygen therapy, non-invasive ventilation, exercise training, pulmonary rehabilitation and lung volume reduction surgery (Barnes P.J., 2000).

A better understanding of the cellular and molecular mechanisms involved in COPD provides new molecular targets for the development of drugs and several classes of new drugs are now under development. Leukotriene B₄ (LT B₄) inhibitors, chemokine inhibitors, TNF- α inhibitors, antioxidants, iNOS inhibitors, and corticosteroids are examples of inflammatory mediator antagonists. Examples of protease inhibitors are neutrophil elastase inhibitors, cathepsin inhibitors, α_1 -antitrypsin, secretory leukoprotease inhibitor and elafin. New anti-inflammatory drugs for COPD are PDE type IV inhibitors, NF- κ B inhibitors, adhesion molecule blockers, IL-10, p38 MAP kinase inhibitors, and PI3-kinase inhibitors (Barnes P.J., 2001).

As described earlier, currently available therapies for T1 inflammatory lung disease are broad acting and directed to the improvement of clinical symptoms and/or to a general reduction of inflammation. There is thus a need for new selective therapeutic strategies which end the relentless progression of said diseases by targeting a key mediator of the underlying mechanism.

Notwithstanding the fact that several potential therapies for T1 mediated inflammatory lung disease have been proposed, no prior art exists revealing that neutralizing IFN γ bioactivity is effective in the treatment of T1 mediated inflammatory lung diseases such as COPD, severe asthma, sarcoidosis, berylliosis, and cystic fibrosis.

The present invention demonstrates that the inflammatory and destructive processes that underlie T1 inflammatory lung diseases can be treated by neutralizing IFN γ .

SUMMARY OF THE INVENTION

It is an object of the present invention to provide methods and compositions for preventing or treating T1 inflammatory lung disease, particularly COPD, emphysema, chronic bronchitis, bronchiolitis, severe asthma, sarcoidosis, berylliosis, and cystic fibrosis.

The present invention provides a method for preventing or treating T1 inflammatory lung disease, said method comprising reducing or neutralizing the bioactivity of IFN γ . Several methods and compositions can be applied for this and are thus part of the current invention.

An aspect of the present invention relates to the use of an IFN γ neutralizing molecule for preventing or treating T1 inflammatory lung disease. More particularly, the present invention relates to the use of an anti-IFN γ antibody for preventing or treating T1 inflammatory lung disease, said antibody preferably being a monoclonal antibody. Furthermore, the present invention relates to the use of a human or a humanized anti-IFN γ antibody for preventing or treating T1 inflammatory lung disease. More specifically, the present invention relates to the use of the anti-IFN γ antibody D9D10, and more particularly a humanized anti-IFN γ antibody D9D10, for preventing or treating T1 inflammatory lung disease.

Another aspect of the present invention relates to the use of immunogenic IFN γ for preventing or treating T1 inflammatory lung disease, and in particular the use of human immunogenic IFN γ .

Accordingly, the present invention also relates to the prevention or treatment of T1 inflammatory lung disease by immunization with a pharmaceutical composition comprising immunogenic IFN γ proteins and/or IFN γ derived (poly)peptides. Several techniques to render IFN γ immunogenic are well known in the art and the present invention allows for all kinds of permutations of the original IFN γ sequence, and all kinds of modifications therein.

Another aspect of the invention relates to the use of the technology of genetic immunization, also known as "DNA vaccination", for preventing or treating T1 inflammatory lung disease.

DETAILED DESCRIPTION OF THE INVENTION

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described. All publications mentioned hereunder are incorporated by reference. Unless mentioned otherwise, the techniques employed herein are standard methodologies well known to one of ordinary skill in the art. The materials, methods and examples are only illustrative and not limiting.

The present invention provides compositions and methods for preventing and treating T1 inflammatory lung disease. Said lung diseases include but are not limited to COPD (comprising emphysema, chronic bronchitis, and bronchiolitis), severe asthma, sarcoidosis, berylliosis and cystic fibrosis. The methods of the invention encompass the neutralization and/or reduction and/or blockade of IFN γ bioactivity. The current invention thus relates to a method for preventing or treating T1 inflammatory lung disease, said method comprising the neutralization of IFN γ bioactivity. Several methods and/or compositions can be used in order to achieve said effect and will be described hereunder.

Administration of IFN γ neutralizing molecules.

A first aspect of the invention is directed to the use of a molecule capable of neutralizing and/or reducing and/or fully inhibiting the bioactivity of IFN γ for preventing or treating T1 inflammatory lung disease. More specifically, the invention relates to the use of an IFN γ neutralizing molecule for the manufacture of a medicament for preventing or treating a T1 inflammatory lung disease.

A "T1 inflammatory lung disease" is characterized by a Type 1 immune response mediated by T helper-1 cells (CD4+) and T cytotoxic-1 cells (CD8+) and by predominant production of interferon gamma (IFN γ), tumor necrosis factor (TNF) and interleukin-2 (IL-2). T1 cytokines evoke cell-mediated immunity characterized by prominent lung tissue infiltration of macrophages, neutrophils and T-cells. Examples of T1 inflammatory lung diseases include, but are not limited to COPD, emphysema, chronic bronchitis, bronchiolitis, severe asthma, sarcoidosis, berylliosis, and cystic fibrosis.

As used herein, the term "molecule" encompasses, but is not limited to, an antibody and fragments thereof, a diabody, a triabody, a tetravalent antibody, a peptide, a low molecular weight non-peptide molecule (also referred to as "small molecules") and a (soluble) IFN γ receptor or fragments thereof, which specifically reduces and/or inhibits IFN γ bioactivity. IFN γ of any species, including humans, is to be considered in this respect.

As used herein, the term "antibody" refers to monoclonal antibodies, polyclonal antibodies, antibodies which are derived from a phage library, humanized antibodies, synthetic antibodies, chimeric antibodies, antibody fragments, single-chain Fv's, or constructs thereof. The term "monoclonal antibody" refers to an antibody composition having a homogeneous antibody population. The term is not intended to be limited by the manner in which it is made. A monoclonal antibody typically displays a single binding affinity for a particular polypeptide with which it immunoreacts. A monoclonal antibody to an epitope of the IFN γ antigen can be prepared by using a technique which provides for the production of antibody molecules by continuous cell lines in culture. These include but are not limited to the hybridoma technique originally described by Kohler and Milstein (1975). Monoclonal antibodies can also be produced in various ways using techniques well understood by those having ordinary skill in the art. Details of these techniques are described in "Antibodies: A Laboratory Manual", Harlow et al.(ed.), Cold Spring Harbor Publications, p. 726 (1988), or are described by Campbell, A.M. ("Monoclonal Antibody Technology Techniques in Biochemistry and Molecular Biology," Elsevier Science Publishers, Amsterdam, The Netherlands (1984)) or by St. Groth et al.(J. Immunol. Methods 35:1-21 (1980)). Monoclonal antibodies of any species, including humans, can be used in this invention. Accordingly, the antibodies according to this embodiment may be

human monoclonal antibodies. Such human monoclonal antibodies may be prepared, for instance, by the generation of hybridomas, derived from immunised transgenic animals, containing large sections of the human immunoglobulin (Ig) gene loci in the germline, integrated by the yeast artificial chromosomal (YAC) technology (Mendez et al., 1997). Also fragments derived from these monoclonal antibodies such as Fab, F(ab)₂ and scFv ("single chain variable fragment"), providing they have retained the original binding properties, form part of the present invention. The present invention thus also relates to the use of an anti-IFN γ antibody for the manufacture of a medicament for preventing or treating T1 inflammatory lung disease, wherein said antibody is a monoclonal or polyclonal antibody, and more particularly a human monoclonal or polyclonal antibody.

As used herein, the term "humanized antibody" means that at least a portion of the framework regions of an immunoglobulin or engineered antibody construct is derived from human immunoglobulin sequences. It should be clear that any method to humanize antibodies or antibody constructs, as for example by variable domain resurfacing as described by Roguska et al. (1994) or CDR grafting or reshaping as reviewed by Hurlle and Gross (1994), can be used.

As used herein, the term "chimeric antibody" refers to an engineered antibody construct comprising variable domains of one species (such as mouse, rat, goat, sheep, cow, llama, or camel variable domains), which may be humanized or not, and constant domains of another species (such as primate or human constant domains) (for review see Hurlle and Gross (1994)). It should be clear that any method known in the art to develop chimeric antibodies or antibody constructs can be used.

As used herein, the term "single chain Fv", also termed scFv, refers to engineered antibodies prepared by isolating the binding domains (both heavy and light chains) of a binding antibody, and supplying a linking moiety which permits preservation of the binding function. This forms, in essence, a radically abbreviated antibody, having only that part of the variable domain necessary for binding the antigen.

Information concerning the generation, design and expression of recombinant antibodies can be found in Mayforth RD, "Designing Antibodies", Academic Press, San Diego (1993).

As used herein, the term "fragment" or "fragments" refers to F(ab), F(ab)₂, Fv, scFv and other fragments which retain the antigen binding function and specificity of the

parent antibody. The methods for producing said fragments are well known to a person skilled in the art and can be found, for example, in *Antibody Engineering*, Oxford University Press, Oxford (1995) (1996) and *Methods in Molecular Biology*, Humana Press, New Jersey (1995). In addition, any construct of an antibody or a fragment is also a subject of the current invention. As used herein, the term "construct" relates to synthetic or recombinant molecules, including but not limited to diabodies, triabodies, tetravalent antibodies, pepta- or hexabodies, and the like, that are derived from an anti-IFN γ antibody. The present invention thus relates to the use of an IFN γ neutralizing molecule for the manufacture of a medicament for preventing or treating a T1 inflammatory lung disease, whereby said molecule is a construct derived from an anti-IFN γ antibody.

As used herein, the term "diabody" relates to two non-covalently-linked scFv's, which then form a so-called diabody, as described in detail by Holliger et al. (1993) and reviewed by Poljak (1994). It should be clear that any method to generate diabodies, as for example described by Holliger et al. (1993), Poljak (1994), and Zhu et al. (1996), can be used.

As used herein, the term "triabody" relates to trivalent constructs comprising 3 scFv's, and thus comprising 3 variable domains, as described by Kortt et al. (1997) and Iliades et al. (1997). A method to generate triabodies is described by Kortt et al. (1997). An example of a triabody is given in WO 99/09055 by Innogenetics N.V.

It should also be clear that the scFv's, chimeric antibodies, diabodies and triabodies described above are not limited to comprise the variable domain of the same antibody but may also comprise variable domains of other anti-IFN γ antibodies which efficiently reduce or neutralize the bioactivity of IFN γ . Furthermore, the diabodies or triabodies described above may also comprise two scFv's of different specificities. For example, the latter diabodies may simultaneously neutralize IFN γ on the one hand and may target another molecule, such as TNF- α , IL-1, IL-2, B7.1 or CD80, B7.2 or CD86, IL-12, IL-4, IL-10, CD40, CD40L, IL-6, complement factor, coagulation factor, fibrinolysis factor, tumour growth factor-beta (TGF- β), transferrin receptor, insulin receptor and prostaglandin E2, or any other molecule, on the other hand.

As used herein the terms "IFN γ neutralizing molecule" or "IFN γ neutralizing antibody" refer to a molecule and an antibody which inhibits or blocks any bioactivity of IFN γ , respectively.

The term "bioactivity" or "biological activity" of IFN γ relates to the antiviral activity (Billiau, 1996), the induction of the expression of MHC-class-II molecules by macrophages and other cell types (Steinman et al., 1980), the stimulation of the production of inflammatory mediators such as TNF- α , IL-1 and NO (Lorsbach et al., 1993), the induction of the expression of adhesion molecules such as ICAM-1 (Dustin et al., 1988) and of important costimulators such as the B7 molecules on professional antigen presenting cells (Freedman et al., 1991), the induction of macrophages to become tumoricidal (Pace et al., 1983), the induction of Ig isotype switching (Snapper and Paul, 1987) or any other known bioactivity of IFN γ . Billiau et al. (1996) describes pathological and/or clinical activity during diseases in which IFN γ is pathogenic. It should be noted that the molecules which neutralize IFN γ as described herein, neutralize at least one bioactivity but not necessarily all bioactivities of IFN γ .

Tests to evaluate the effect of anti-IFN γ molecules or antibodies (i.e. IFN γ neutralizing molecules or antibodies resp.) on the bioactivity of IFN γ are available and well known to the skilled person. Examples of said tests are, but not limited to, "inhibition of MHCII-induction" and/or "inhibition of anti-viral activity". In the first mentioned test, the effect of IFN γ on the induction of MHC class II expression on keratinocytes is examined. For this, primary keratinocytes are cultured with two concentrations of IFN γ (100 U/ml and 200 U/ml) for 24 and 48 hours. After culture, cells are collected and the expression of MHC class II antigen on the activated keratinocytes is measured by FACS-scan after staining (30 minutes at 4°C) of the cells with a PE-labelled anti-MHC-class II mAb. In addition, the effect of an anti-IFN γ molecule on the IFN γ -induced MHC-Class II expression on keratinocytes is examined. In this experiment, primary keratinocytes are cultured with IFN γ (100 U/ml) in the presence or absence of different concentrations of anti-IFN γ molecules or antibodies for 48 hours. IFN γ is preincubated with anti-IFN γ molecules or antibodies for 1 hour at 37°C before adding to the keratinocytes. After culture, cells are collected and the expression of MHC-Class II on these activated keratinocytes is measured. For this, keratinocytes are incubated (30 minutes at 4°C) with a PE-labelled anti-MHC-ClassII mAb (Becton Dickinson), washed twice with PBS and fixed. The MHC-Class II expression is further analysed on a FACS-scan. Analogous to the described test, the effect of IFN γ on the induction of MHC-class II expression on B cells can be examined. Also other experiments known to those skilled in the art

can be performed in order to evaluate the neutralization capacity of anti-IFN γ molecules (incl. antibodies).

For the second test, whereby neutralization of the antiviral activity of IFN γ is measured, serial dilutions of samples (anti-IFN γ molecules or antibodies) are prepared in microtiter plates. IFN γ is added to each well in a final concentration of 5 antiviral protection Units/ml, as tested on A549 cells. The mixtures are incubated for 4 h at 37°C and 25000 A549 cells are added to each well. After an incubation period of 24 at 37°C in a CO₂ incubator, 25 μ l of 8×10^5 PFU EMC virus/ml is added to the cultures for at least 24h. As soon as virus-infected control cultures reach 100% cell destruction, a crystal violet staining is performed in order to quantify surviving cells. The neutralization capacity of the anti-IFN γ molecules or antibodies can be defined for instance, as the concentration of the molecule or antibody needed to neutralize 95% of the antiviral activity of 5U/ml IFN γ . The neutralization potency of the anti-IFN γ molecules or antibodies is then determined. Further assays to evaluate the effect of anti-IFN γ molecules (incl. antibodies) on the bioactivity of IFN γ are described by e.g. Lewis J.A. (1995), Kim Y. & Son K. (1996) and by Maeger A. (2002).

The term "prevention" or "treatment" as used herein refers to either (i) the prevention of the disease of interest (prophylaxis), or (ii) the reduction or elimination of symptoms, exacerbations or the disease of interest (therapy), or (iii) any process, action, application, therapy, or the like, wherein a mammal, including a primate and more specifically a human being, is subject to medical aid with the object of improving the mammal's condition, directly or indirectly.

The present invention thus relates to a method for preventing or treating T1 inflammatory lung disease comprising administering to a patient a pharmaceutically effective amount of an IFN γ neutralizing molecule. More specifically, the present invention relates to a method for preventing or treating T1 inflammatory lung disease comprising administering to a patient a pharmaceutically effective amount of an anti-IFN γ antibody. According to a specific embodiment, the antibody is the monoclonal antibody D9D10H3G5 produced by the hybridoma deposited on August 28, 2001, under the Accession No. DSM ACC2521, with the DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Germany. Said monoclonal antibody D9D10H3G5 will be further abbreviated throughout the specification and the claims as D9D10. More specifically,

the present invention relates to a method for preventing or treating T1 inflammatory lung disease comprising administering to a patient a pharmaceutically effective amount of an anti-IFN γ antibody D9D10 or a fragment thereof. Furthermore, the present invention relates to the use of an anti-IFN γ antibody for the manufacture of a medicament for preventing or treating T1 inflammatory lung disease, said antibody preferably being a monoclonal antibody (e.g. D9D10 as described herein) or a humanized monoclonal antibody (e.g. humanized D9D10 as described herein or as described in WO99/09055 by Innogenetics N.V.).

Differently produced antibodies recognizing the same epitopes as the antibody D9D10, as well as antibodies immunologically competing with the antibody D9D10 for the binding on IFN γ are also part of the invention. Therefore, according to a further embodiment, the present invention relates to the use of an anti-IFN γ antibody or a fragment thereof, for preventing or treating T1 inflammatory lung disease, whereby said antibody is characterized by its ability to immunologically compete with the antibody D9D10 for the binding on IFN γ . As used herein, the term "to bind in an equivalent way" or "immunologically competing" means that these antibodies inhibit the binding of D9D10 to IFN γ and that these antibodies neutralize the bioactivity of IFN γ . Preferred methods for determining antibody specificity and affinity by competitive inhibition, e.g. solid phase ELISA, can be found in Harlow et al. (1988), Colligan et al. (1992, 1993), Ausubel et al. (1987, 1992, 1993), and Muller R. (1993), and in Karlsson et al. (1991) and Malmqvist M. (1999).

As used herein, the term "pharmaceutical composition" or "composition" or "medicament" refers to any composition comprising a molecule, including an antibody or fragment thereof, which specifically neutralizes IFN γ , preferably in the presence of a pharmaceutically acceptable carrier or excipient. The present invention thus also relates to the use of a pharmaceutical composition comprising at least one IFN γ neutralizing molecule and an acceptable carrier for preventing or treating T1 inflammatory lung disease. More preferably, said composition comprises the antibody D9D10 or a humanized D9D10 antibody and a pharmaceutically acceptable carrier. Further, said composition optionally comprises other drugs or other antibodies, antibody derivatives or constructs. With regard to COPD, emphysema, chronic bronchitis and chronic obstructive bronchiolitis, examples of such other drugs or other

antibodies, antibody derivatives or constructs are, but are not limited to: bronchodilators, corticosteroids, theophylline, antibiotics, Leukotriene B₄ (LT B₄) inhibitors, chemokine inhibitors, TNF- α inhibitors, anti-IL-8, antioxidants, iNOS inhibitors, neutrophil elastase inhibitors, cathepsin inhibitors, α_1 -antitrypsin, secretory leukoprotease inhibitors, elafin, PDE type IV inhibitors, NF- κ B inhibitors, adhesion molecule blockers, IL-10, p38 MAP kinase inhibitors, PI3-kinase inhibitors and TNF-tip peptides as described in WO 00/09149; with regard to severe asthma: β_2 -agonists, anticholinergics, corticosteroids (e.g. prednisone); with regard to sarcoidosis: corticosteroids; cytotoxic agents, immunomodulators (e.g. chloroquine, hydroxychloroquine and TNF inhibitors such as pentoxifylline and thalidomide) and the antileprosy drugs clofazimine and minocycline; with regard to berylliosis: corticosteroids; with regard to cystic fibrosis: antimicrobial agents, corticosteroids and non-steroidal anti-inflammatory drugs such as ibuprofen.

It should also be clear that any possible mixture of any IFN γ neutralizing molecule, antibody or composition described in the specification may be part of the above-indicated pharmaceutical composition. The proportion and nature of said pharmaceutical compositions are determined by the solubility and chemical properties of the selected compound, the chosen route of administration, and standard pharmaceutical practice.

The IFN γ neutralizing molecule, antibody or a fragment thereof, and more preferred the monoclonal antibody D9D10 or a humanized D9D10 antibody, or a fragment or construct thereof, may thus be administered or delivered in the form of any suitable composition as described in the specification by any suitable method of administration within the knowledge of one skilled in the art.

As used herein, the term "pharmaceutically acceptable carrier or excipient", whereby the term carrier and excipient are used interchangeably, refers to a diluent, adjuvant, or vehicle with which the therapeutic molecule is administered. It includes any and all solvents, dispersion media, aqueous solutions, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, the use thereof in pharmaceutical compositions is contemplated. Supplementary active

ingredients can also be incorporated into the compositions of the invention. A composition is said to be "pharmacologically acceptable" if its administration can be tolerated by the recipient.

Immunization.

Another method to neutralize the bioactivity of IFN γ is by immunization.

There is an increased focus on methods of instructing the recipient's own immune system to generate endogenous antibodies of the appropriate specificity by means of immunization. However, mammals do not generally have high-titre antibodies against self-proteins in serum because of homeostatic tolerance mechanisms that prevent their formation (autotolerance).

It has been shown (by Dalum L. et al., 1996) that potentially self-reactive B lymphocytes recognizing self-proteins are physiologically present in normal individuals. However, in order for these B lymphocytes to be induced to actually produce antibodies reactive with the relevant self-proteins, assistance is needed from cytokine producing T helper lymphocytes (Th-cells or Th-lymphocytes). Normally this help is not provided because T lymphocytes in general do not recognize T-cell epitopes derived from self-proteins when presented by antigen presenting cells (APCs) such as dendritic cells, macrophages and B cells. However, by providing an element of "foreignness" in a self-protein, T-cells recognizing the foreign element are activated upon recognizing the foreign epitope on an APC (such as, initially, a mononuclear cell). Polyclonal B lymphocytes (which are also specialised APCs) capable of recognizing self-epitopes on the modified self-protein, also internalize the antigen and subsequently present the foreign T-cell epitope(s) thereof, and the activated T lymphocytes subsequently provide cytokine help to these self-reactive polyclonal B lymphocytes. Since the antibodies produced by these polyclonal B lymphocytes are reactive with different epitopes on the modified polypeptide, including those which are also present in the native polypeptide, an antibody cross-reactive with the non-modified self-protein is induced. In conclusion, the T lymphocytes can be led to act as if the population of polyclonal B lymphocytes have recognized an entirely foreign antigen, whereas in fact only the added or inserted

epitope(s) is/are foreign to the host. In this way, antibodies capable of cross-reacting with non-modified self-antigens are induced.

Different kinds of immunization approaches that are able to break B-cell tolerance and circumvent antibody tolerance mechanisms without inducing auto-antibody-mediated pathology and toxicology are well known by the skilled person and are outlined hereunder.

Accordingly, the present invention relates to the prevention or treatment of T1 inflammatory lung disease by immunization with a pharmaceutical composition comprising immunogenic IFN γ proteins and/or IFN γ derived (poly)peptides. These IFN γ -related compounds are capable of raising auto-antibodies and/or T cells when administered in vivo. The auto-antibodies are able to neutralize and inhibit the biological activities of endogenously produced IFN γ . The present invention thus relates to the use of immunogenic IFN γ or IFN γ derived (poly)peptides for the manufacture of a medicament for preventing or treating a T1 inflammatory lung disease.

"Immunization" means that a substance or composition of matter exhibits or induces an immune response resulting in endogenous antibody production concomitant with or without T-cell help. Endogeneous antibody production against IFN γ can be measured using standard techniques, e.g. by ELISA.

The term "immunogen" is intended to denote a substance which is capable of inducing an immune response in a certain mammals, including primates and more specifically humans. It will therefore be understood that autologous IFN γ is not an immunogen in the autologous host under non-pathogenic conditions. It is necessary to use either a strong adjuvant and/or to co-present foreign T helper epitopes with the autologous IFN γ in order to mount an immune response against autologous IFN γ and in such a case the "immunogen" is the composition of matter which is capable of breaking autotolerance. Other immunogens described in the current invention include but are not limited to modified IFN γ , non-self IFN γ (i.e. IFN γ from another species) and antigen presenting cells loaded with IFN γ . The term "immunogenic IFN γ " thus specifically relates to:

- modified IFN γ and/or,
- non-self IFN γ derived from another species (mammalian or other) administered with or without an adjuvant and/or,
- autologous IFN γ co-presented or in combination with an adjuvant and/or,
- autologous antigen presenting cells, such as dendritic cells, loaded with IFN γ .

The expressions "IFN γ ", "IFN γ protein" and "IFN γ polypeptide", which are used interchangeably, refer to a family of polypeptide molecules that include human IFN γ from natural sources, synthetically produced in vitro, or obtained by genetic manipulation including methods of recombinant DNA technology. The amino acid sequence variants preferably share at least about 65% sequence homology, more preferably at least about 75% sequence homology, even more preferably at least about 85% sequence homology, most preferably at least about 90% sequence homology with any domain, and preferably with the receptor binding domain(s) of the native human IFN γ amino acid sequence. Several databases and tools for determining amino acid homology are known by the person skilled in the art, e.g. BLAST®, and are described by Gish W. & States D.J. (1993) and in "Bioinformatics: sequence and genome analysis", Mount (ed.), Cold Spring Harbor Laboratory Press (2001). The definition specifically covers variously glycosylated and unglycosylated forms of native human IFN γ and of its amino acid sequence variants.

A "modified IFN γ " is an IFN γ protein or IFN γ derived (poly)peptide which has been subjected to changes in its primary structure. Such a change can e.g. be in the form of fusion or conjugation of an IFN γ polypeptide to a suitable fusion partner (i.e. a change in primary structure exclusively involving C-and/or N-terminal additions of amino acid residues) and/or it can be in the form of insertions and/or deletions and/or substitutions in the amino acid sequence of IFN γ . It should also be noted that the IFN γ protein can become immunogenic due to certain modifications resulting from the recombinant production process, isolation, handling, or storage of the protein. According to a preferred embodiment of the present invention the IFN γ protein is rendered immunogenic by its modification.

One technique involves chemically cross-linking of the IFN γ self-protein (or (poly)peptide(s) derived from it) to a highly immunogenic non-self carrier protein such as, but not limited to, keyhole limpet haemocyanin (Kim, 2001), ovalbumin (Richard, 2000), tetanus toxoid, detoxified diphtheria or cholera toxin (Talwar, 1999), bacterial outer membrane proteins (Gonzalez, 2000), E.coli enterotoxin (Lowenadler, 1994), heat shock proteins (Udono, 1993), and viral-like particles (Chackerian, 2002; Storni, 2002). The carrier protein contains several different foreign T-cell epitopes needed to trigger the activation of T-cells which in their turn provide help to B-cells that produce antibodies specifically directed against the conjugated IFN γ self-antigen. Instead of whole carrier proteins, the chemical cross-linking of foreign MHC Class II restricted T-cell epitopes may also be efficient for the induction of auto-immune responses against IFN γ (Sad, 1992). The current invention thus relates to the use of immunogenic IFN γ for the manufacture of a medicament for preventing or treating a T1 inflammatory lung disease, whereby the IFN γ protein is modified to immunogenic IFN γ by crosslinking IFN γ to an immunogenic non-self carrier protein. Furthermore, the current invention also relates to the use of immunogenic IFN γ for the manufacture of a medicament for preventing or treating a T1 inflammatory lung disease, whereby the IFN γ protein is modified to immunogenic IFN γ by chemical cross-linking of one or more foreign MHC Class II restricted T-cell epitopes.

A variant on the carrier protein technique involves the construction of a gene encoding a fusion protein comprising both carrier protein or foreign T-cell epitope(s) and the IFN γ self-protein or B-cell epitope(s) derived therefrom. The fusion protein may be expressed in a suitable host cell in vitro, the gene product purified and then delivered as a conventional pharmaceutical composition co-presented with or without adjuvant. The current invention thus also relates to the use of a fusion protein for the manufacture of a medicament for preventing or treating a T1 inflammatory lung disease, whereby the fusion protein comprises a carrier protein or at least one foreign T-cell epitope and the IFN γ self-protein or B-cell epitope(s) derived therefrom. Alternatively, the fusion gene may be administered directly as part of a nucleic acid vaccine.

The foreign T-cell epitope(s) can not only be conjugated to the amino- or carboxy-terminal end of the human IFN γ self-protein but can alternatively be inserted into the human IFN γ protein either at random or at sites predicted not to interfere with the general conformational structure. As a result, T-cell help arises either from this epitope(s) or from junctional sequences.

A more refined approach has been described by Dalum and colleagues wherein a small part of target molecule is substituted by a single MHC Class II restricted T cell epitope (Dalum L, 1999). The same approach can be used to introduce multiple epitopes by substitution. The substituted epitope is supposed not to interfere with the folding of the protein resulting in the adoption of a fully native conformation identical to the one found in the native endogenous self-protein. Therefore, it is highly probable that antibodies recognizing the immunogen will also recognize and at best neutralize the endogenous self-protein.

One preferred version of this embodiment is the technique described in WO 95/05849 by Elsner Henrik et al., which discloses a method for down-regulating self-proteins by immunizing with analogues of the self-proteins wherein a number of amino acid sequence(s) has been substituted with a corresponding number of amino acid sequence(s) which each comprise a foreign immunodominant T-cell epitope, while at the same time maintaining the overall tertiary structure of the self-protein in the analogue. For the purposes of the present invention it is, however, sufficient if the modification (be it an amino acid insertion, addition, deletion or substitution) gives rise to a foreign T-cell epitope and at the same time preserves a substantial number of the B-cell epitopes in IFN γ . The present invention thus relates to the use of immunogenic IFN γ for the manufacture of a medicament for preventing or treating a T1 inflammatory lung disease, wherein the IFN γ protein is modified to immunogenic IFN γ by introducing at least one foreign T-cell epitope into IFN γ by insertion, substitution, addition and/or conjugation. Said foreign T-cell epitope can be any natural or synthetic T-cell epitope as described herein.

A "foreign T-cell epitope" is a peptide which is able to bind to an MHC molecule and stimulates T-cells in an animal species. Preferred foreign epitopes are "promiscuous" epitopes, i.e. epitopes which bind to a substantial fraction of polymorphic MHC class

II molecules in an animal species. A term which is often used interchangeably in the art is the term "universal T-cell epitopes" for this kind of epitope.

A number of naturally occurring "promiscuous" T-cell epitopes exist which are active in a large proportion of individuals of an animal species, including human, and these are preferably introduced in the IFN γ composition thereby reducing the need for a very large number of different analogues in the same composition.

The promiscuous epitope can according to the invention be a naturally occurring human T-cell epitope such as epitopes from tetanus toxoid (e. g. the P2 and P30 epitopes in WO 00/20027), diphtheria toxoid, influenza virus hemagglutinin (HA), and *P. falciparum* CS antigen.

Over the years a number of other promiscuous T-cell epitopes have been identified. In particular, peptides capable of binding a large proportion of HLA-DR molecules encoded by the different HLA-DR alleles have been identified and these are all possible T-cell epitopes to be introduced in analogues used according to the present invention. Additional epitopes are discussed in the following references: Kilgus et al., 1991; Contreras et al., 1998; Doolan et al., 2000; Launois et al., 1994; Mustafa et al., 2000; Fernando et al., 1995; Gaudebout et al., 1997; Friedl-Hayek et al., 1999; Kobayashi et al., 2000. All epitopes listed in these references are relevant as candidate epitopes to be used in the present invention, as are epitopes which share common motifs with these.

Alternatively, the epitope can be any synthetic or artificial T-cell epitope which is capable of binding a large proportion of haplotypes. In this context the pan DR epitope peptides ("PADRE") described in WO 95/07707 and in the corresponding paper by Alexander J. et al. (1994) are interesting candidates for epitopes to be used according to the present invention.

Another mechanism provides for the generation of a multiplicity of potential T-cell epitopes, yet simultaneously retains the target molecule in a conformation close to the native form (Ciapponi, 1997). These properties can be achieved by rendering one or several mutations in a self-protein to produce a sequence at those points which can be found in an analogous protein from a second mammalian species. Mutations can be applied to consecutive amino acids or locally dispersed amino acids.

The current invention relates to the use of immunogenic IFN γ for the manufacture of a medicament for preventing or treating a T1 inflammatory lung disease, whereby the IFN γ protein is modified to immunogenic IFN γ by rendering one or several mutations in the IFN γ self-protein. Such mutations may in effect produce a sequence at those points which can be found in an analogous protein from a second mammalian species. Alternatively, B-cell epitopes of IFN γ from a first mammalian species may be grafted (e.g. by insertion or substitution) into the framework of a protein from a second mammalian species such that the modified protein is able to raise in the first species an immunogenic response directed to the natural IFN γ protein from which the B-cell epitopes are derived. A specific embodiment of the current invention thus relates to the use of immunogenic IFN γ for the manufacture of a medicament for preventing or treating a T1 inflammatory lung disease, whereby a protein from a second mammalian species is modified to an immunogen by grafting B-cell epitopes from an IFN γ protein from a first mammalian species into the frame work of protein from the second mammalian species. Preferably, the first mammalian species is human.

Auto-antibodies in humans can also be induced by immunization with a non-modified analogous protein from a second mammalian species (Vernersson, 2002). Antibodies directed against the analogous protein are suspected to cross-react with the human self-protein resulting in the neutralization of its biological activity. The invention thus also relates to the use of immunogenic IFN γ for the manufacture of a medicament for preventing or treating a T1 inflammatory lung disease, whereby the immunogenic IFN γ is a non-self IFN γ derived from a second mammalian species.

Anti-human IFN γ auto-antibodies in humans can also be achieved by immunization with autologous antigen presenting cells (APCs) ex vivo loaded with human IFN γ (Li, 2002). The invention thus also relates to the use of at least one autologous antigen presenting cell loaded with IFN γ for the manufacture of a medicament for preventing or treating a T1 inflammatory lung disease. More particularly, the invention relates to the use of at least one autologous dendritic cell loaded with IFN γ for the manufacture of a medicament for preventing or treating a T1 inflammatory lung disease

Thus, given the general functional restraints on the immunogenicity of the constructs, the invention allows for all kinds of permutations of the original IFN γ sequence, and all kinds of modifications therein.

Some of the IFN γ proteins of the pharmaceutical composition are sufficiently immunogenic, but for some the immune response will be enhanced if the composition further comprises an adjuvant substance. It is especially preferred to use an adjuvant which can be demonstrated to facilitate breaking of the autotolerance to autoantigens. Various methods of achieving adjuvant effect for the composition are known. General principles and methods are detailed in "The Theory and Practical Application of Adjuvants", 1995, Duncan E. S. Stewart-Tull (ed.), John Wiley & Sons Ltd., and also in "Vaccines : New Generation Immunological Adjuvants", 1995, Gregoriadis G et al. (eds.), Plenum Press, New York.

Preferred adjuvants facilitate uptake of the molecules by APCs, such as dendritic cells, and activate these. Non-limiting examples are selected from the group consisting of an immune targeting adjuvant; an immune modulating adjuvant such as a toxin, a cytokine, and a mycobacterial derivative; an oil formulation ; a polymer; a micelle forming adjuvant; a saponin; an immunostimulating complex matrix (ISCOM matrix); a particle; DDA aluminium adjuvants; DNA adjuvants; γ -inulin; and an encapsulating adjuvant. Details relating to composition and use of immunostimulating complexes can e.g. be found in the above-mentioned text-books dealing with adjuvants, but also Morein B. et al., 1995, as well as Barr I.G. and Mitchell G.F., 1996, provide useful instructions for the preparation of complete immunostimulating complexes. The current invention thus also relates to the use of a pharmaceutical composition comprising immunogenic IFN γ and an adjuvant for treating T1 inflammatory lung disease. More specifically, the current invention relates to the use of a pharmaceutical composition comprising autologous IFN γ and an adjuvant for treating T1 inflammatory lung disease.

Another approach of immunization includes the technology of nucleic acid immunization, also known as "genetic immunization", "gene immunization" and "DNA vaccination". The present invention thus also relates to the use of a DNA

vaccine for preventing or treating T1 inflammatory lung disease.

In this embodiment, the introduced nucleic acid encoding the modified IFN γ protein as described herein is preferably DNA which can be in the form of naked DNA, DNA formulated with charged or uncharged lipids, DNA formulated in liposomes, emulsified DNA, DNA included in a viral vector, DNA formulated with a transfection-facilitating protein or polypeptide, DNA formulated with a targeting protein or polypeptide, DNA formulated with calcium precipitating agents, DNA coupled to an inert carrier molecule, and DNA formulated with an adjuvant. In this context it is noted that practically all considerations pertaining to the use of adjuvants in traditional vaccine formulation apply to the formulation of DNA vaccines. Hence, all disclosures herein which relate to use of adjuvants in the context of protein or (poly)peptide based pharmaceutical compositions apply *mutatis mutandis* to their use in nucleic acid vaccination technology. The same holds true for other considerations relating to formulation and mode and route of administration and, hence, also these considerations discussed herein in connection with a traditional pharmaceutical composition apply *mutatis mutandis* to their use in nucleic acid vaccination technology.

Accordingly, the present invention also relates to a method of preventing or treating T1 inflammatory lung disease by immunization with a pharmaceutical composition comprising a nucleic acid sequence encoding an immunogenic IFN γ protein and/or IFN γ -derived (poly)peptide. More specifically, the present invention relates to the use of a nucleic acid sequence encoding an immunogenic IFN γ protein and/or IFN γ -derived (poly)peptide for the manufacture of a medicament for preventing or treating a T1 inflammatory lung disease.

Under normal circumstances, the nucleic acid of the vaccine is introduced in the form of a vector wherein expression is under control of a viral promoter.

Therefore, also provided are an expression vector which comprises a polynucleotide of the herein described proteins or peptides and which is capable of expressing the respective proteins or peptides, a host cell comprising the expression vector and a method of producing and purifying herein described proteins or peptides, pharmaceutical compositions comprising the herein described proteins or peptides and a pharmaceutically acceptable carrier and/or adjuvants.

Detailed disclosures relating to the formulation and use of nucleic acid vaccines are available, e.g. by Donnelly J.J. et al, 1997 and 1997a.

Administration

The molecule, protein, composition or agent of the current invention may be administered in any manner which is medically acceptable. In addition, it can at any time be administered together, simultaneously or sequentially, with another separate substance, molecule, antibody, composition, or a pharmaceutically and immunologically acceptable carrier and/or vehicle and/or an adjuvant.

Any of the conventional methods for administration of a pharmaceutical composition are applicable e.g. parenterally, orally or by either subcutaneous, intradermal, subdermal, intravenous, intraarterial or intramuscular injection. Other modes of administration include suppositories and, in some cases, buccal, sublingual, intraperitoneal, intravaginal, anal, and intracranial applications. Depending on the specific circumstances, local or systemic administration may be desirable.

One skilled in the art of preparing formulations can readily select the proper form and mode of administration depending upon the particular characteristics of the molecule, protein, composition or agent selected, the disease state to be treated, the stage of the disease, and other relevant circumstances.

According to the specific case, the "pharmaceutically effective amount" or "amount effective" is one that is sufficient to produce the desired effect. This can be monitored using several end-points known to those skilled in the art such as, but not limited to, mortality, morbidity, and the like. According to the specific case, the pharmaceutically effective amount should be determined as being the amount sufficient to cure the recipient in need of treatment, to prevent or at least to partially reduce or halt the disease or injury and its complications. The term "recipient" is intended to include living organisms, e.g. mammals, primates, and more specifically humans. Amounts effective for such use will depend on the severity of the disease and the general state of the recipient's health. As such, dosage of the administered molecule, protein, composition or agent will vary depending upon such factors as the recipient's age, weight, height, sex, general medical condition, previous medical

history, concurrent treatment with other pharmaceuticals, etc. Administration can be as a single dose or repeated doses one or more times after a specific period. When administering by injection, the administration may be by continuous injection, or by single or multiple boluses. Typically, such compositions are prepared as injectables either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in liquid prior to injection may also be prepared. The preparation may also be emulsified.

The active ingredient of the pharmaceutical composition is often mixed with excipients or carriers which are pharmaceutically acceptable and compatible with the active ingredient. Such excipients are inherently nontoxic and nontherapeutic.

Examples of such excipients are water, saline, glycerol, ethanol, Ringer's solution, dextrose solution and Hank's solution, and/or combinations thereof. Nonaqueous excipients such as fixed oils and ethyl oleate may also be used. A preferred excipient is 5% dextrose in saline. The excipient may contain minor amounts of additives such as substances that enhance isotonicity and chemical stability, including buffers and preservatives. In addition, if desired, the composition may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, or adjuvants which enhance the effectiveness of the composition. Examples of adjuvants are sterile diluents such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylene diaminetetraacetic acid; buffers such as acetates, citrates or phosphates, and agents for the adjustment of tonicity such as sodium chloride or dextrose.

LEGEND TO THE FIGURE

Figure 1: TNF- α levels were measured in the serum of the animals after 6 weeks of smoking. Sera from 3 placebo-treated animals and from 3 animals from the anti-IFN γ treated group were tested. TNF- α was measured using an in house developed ELISA. The detection limit is 4pg/ml. As shown in this figure, treatment of the smoking animals with anti-IFN γ inhibits the TNF- α production.

EXAMPLES

Example 1:

Measurement of IFN γ in lungs of patients with T1 inflammation

Immunological inflammation may be of a Type 1 with predominance of interleukin 2 (IL-2) and interferon gamma (IFN γ) production or a Type 2 characterized by predominant IL-4 and IL-5 production. Presence of IFN γ in the lungs of patients with T1 inflammatory lung diseases can be measured in different ways as outlined hereunder.

One method is the measurement of cytokine levels in spontaneously produced and/or induced sputum. Sputum is defined as expectorations of fluid, cells and solutes that are present in the lining fluid of the upper bronchial tree. Spontaneous sputum can be obtained in a simple and non-invasive way whilst induced sputum is induced by exposure of individuals to a nebulised saline solution (Out et al. 2001). Protease inhibitors can be added shortly after isolation.

The gel and sol phase in sputum can be separated from each other by means of ultracentrifugation (50.000g; 4°C). Otherwise, mucus can be homogenized by treatment with dithiotreitol (DTT) (Karpati et al., 2000). Additional treatment by DNase to degrade any DNA present or treatment with trypsin-EDTA may be necessary. Further low speed centrifugation (300g) of homogenised sputum allows the separation of supernatant from the cellular compartment. Supernatant can be used as such or concentrated and eventually stored at -70°C until assessment of cytokine levels.

BAL fluid is obtained by standard lavage protocol consisting of infusing 20 ml aliquots of sterile saline solution through an aspiration port followed by lavage collection through same. This procedure is repeated 5 times.

Sputum and/or BAL is collected from severe COPD patients (FEV1 < 30% predicted according to the GOLD criteria), moderate COPD patients (30% \leq FEV1 < 80% predicted according to the GOLD criteria), pack-year matched healthy smokers

(normal FEV1 and no chronic symptoms of cough or sputum production) and control non-smokers. Patients are well characterized: age, gender, pack-years smoking history, lung function test, acute exacerbations, inflammation.

Fluid phase of sputum or BAL is evaluated for presence of IFN γ by means of the Enzyme Linked Immunosorbent Assay (ELISA) method. A proteolytic environment like sputum and the treatment of sputum with reducing agents (DTT) may affect antigenic determinants as well as the characteristics of the antibodies and thus the performance of the immunoassays. Therefore, a particular assay has to be validated for its application in sputum. Commercially available kits from Medgenix or R&D Systems or Amersham Biosciences can be used to determine IFN γ immunoreactivities (Keatings, 2002).

Cellular fractions can be assessed for the presence of relative proportions of different cell types by Wright stain on cytopsin slides. Cellular fractions can further be analysed for IFN γ production by immunohistochemistry on cytopsin slides or RT-PCR on total cell RNA or by FACS analysis of intracellular cytokines.

The leakage of plasma proteins (e.g. albumin, fibrinogen) from the blood into the airspace may be considered as an overall marker for inflammation.

Presence of IFN γ can also be assessed in biopsies of smokers, non-smokers or ex-smokers with mild COPD or in lung fragments obtained by lobectomy of severe COPD patients. IFN γ can be evaluated by immunohistochemistry or RT-PCR.

Example 2:

Effect of blocking Ifn-gamma, by anti-mouse Ifn-gamma Mab, on cigarette smoke induced emphysema in mice

INTRODUCTION

Chronic Obstructive Pulmonary Disease (COPD) is characterized by the progressive development of a not fully reversible airflow limitation (Pauwels et al., 2001). The airflow limitation is due to a variable mixture of respiratory bronchiolitis and emphysema. A major risk factor for COPD is cigarette smoking. Inflammation of the airways and the lungs is thought to play a major role in the pathogenesis of COPD. Human studies clearly illustrate a smoking-induced inflammatory response, comprised of neutrophils, macrophages, dendritic cells, eosinophils, CD4⁺ and CD8⁺

T-lymphocytes in smokers' airways and lung parenchyma (Retamales et al., 2001; Casolaro et al., 1988). The exact role of individual inflammatory cells and mediators is unknown at the moment.

Wang et al (2000) demonstrated that interferon gamma (IFN-gamma) causes emphysema with alveolar enlargement, enhanced lung volumes, enhanced pulmonary compliance, and macrophage- and neutrophil-rich inflammation when inducibly targeted, in a transgenic fashion, to the adult murine lung. Prominent protease and antiprotease alterations were also noted in these mice. They included the induction and activation of matrix metalloproteinase (MMP)-12 and cathepsins B, H, D, S, and L, the elaboration of MMP-9, and the selective inhibition of secretory leukocyte proteinase inhibitor.

The aim of the current study is primary to investigate the effect of blocking IFN-gamma on the development of emphysema in mice exposed for six months to cigarette smoke. A secondary objective is to study the effect of anti-mouse IFN-gamma mAb on the cigarette smoke induced pulmonary inflammation.

DESIGN OF THE STUDY

Three groups of 15 mice are followed for 24 weeks.

Group I is exposed to room air and treated twice a week (Monday and Thursday) intraperitoneally with vehicle.

Group II is exposed to cigarette smoke 5 days a week and treated twice a week (Monday and Thursday) intraperitoneally with vehicle.

Group III is exposed to cigarette smoke 5 days a week and treated twice a week (Monday and Thursday) intraperitoneally with 100µg anti-mouse IFN-gamma mAb, F3.

Blood samples (300-400µl) are taken weekly (Thursday) from the ocular sinus. Serum is prepared from each sample. Serum concentrations of anti-IFN-gamma mAb F3, IFN-gamma, IL-6, TNF alfa, IL-8, IL-16, different chemokines and other parameters, are analyzed using ELISA technology. NO was measured using Colormetric Nitric Oxide detection kit.

Immediately following the last exposure, animals are sacrificed and emphysema and lung inflammation are quantified using the following parameters:

Emphysema: Mean Linear Intercept, Mean Alveolar Surface, Mean number of alveolar wall breaks/alveolar space.

Inflammation:

BAL fluid: total number of cells, differential cell count for macrophages/monocytes, neutrophils, eosinophils and lymphocytes.

Lung tissue (single cell suspension): total number of cells, macrophages, dendritic cells, lymphocytes (CD3+), CD4+ lymphocytes, CD8+ lymphocytes, activated CD4+ lymphocytes, activated CD8+ lymphocytes.

METHODS

Animals

Male C57Bl/6 mice, 10 wks old (at the start of the experiment, i.e. pre-bleeding), are purchased from Harlan (Zeist, The Netherlands). The mice are kept in standard animal research facilities and receive food and water ad libitum. Mice are kept in groups of 6. The local Ethics Committee approved all in vivo manipulations.

Experimental design

Groups of 5 mice are exposed to the tobacco smoke of 5 cigarettes (Reference Cigarette 1R3, University of Kentucky, Lexington) per exposure. There are 4 exposures a day with a 30 minutes smoke-free interval between each exposure. The animals smoke 5 days a week for up to 24 weeks. The control group is exposed to air. At week 24, mice are sacrificed and inflammatory parameters are examined in bronchoalveolar lavage (BAL) fluid and lung single cell suspension. Besides, histological evaluation of lung parenchyma is performed.

Tobacco smoke chamber

Mice are exposed to cigarette smoke, with the use of a smoking apparatus (S. Shapiro, Washington University Medical Center, USA) with the chamber adapted for a group of 6 mice (chamber volume of 7500 cm³). An optimal smoke:air ratio of 1:12 is obtained by injecting smoke and pressurized air at a flow rate of 200 ml/min and 2.4 L/min respectively.

Bronchoalveolar lavage

Animals are sacrificed after i.p. injection of an overdose of pentobarbital (Sanofi, Libourne, France) and the trachea is surgically exposed and cannulated to perform bronchoalveolar lavage. 1 ml of Hank's balanced salt solution (HBSS) (GIBCO BRL), free of ionized calcium and magnesium, and supplemented with 0.05 mM

EDTA (Sigma) is instilled 3 times via the tracheal cannula and recovered by gentle manual aspiration. The recovered bronchoalveolar lavage fluid (BAL) is centrifuged, the cell pellet is washed twice and finally resuspended in 1 ml of HBSS. A total cell count is performed in a Bürcker chamber and the differential cell counts (on at least 400 cells) are performed on cytocentrifuged preparations after May-Grünwald-Giemsa staining. Flow cytometric analysis of BAL-cells is also performed (see below).

The supernatant of the BAL is stored at -80°C for measurement of IFN-gamma.

Buffers and media for preparation of single cell suspensions and immunofluorescent labeling

Digestion medium consists of RPMI 1640 supplemented with 2 mM L-glutamine, 10 $\mu\text{g/ml}$ streptomycin, 100 U/ml penicillin, 5 % FCS, 0.001 % β -mercaptoethanol (all from GIBCO BRL), 1 mg/ml collagenase type 2 (Worthington Biochemical Corp.), and 0.02 mg/ml DNase I (grade II from bovine pancreas; Boehringer). FACS-EDTA buffer contains PBS (GIBCO BRL) without Ca^{2+} or Mg^{2+} , 0.1 % azide, 5 % EDTA-treated FCS, and 5 mM EDTA. EDTA-treated FCS is prepared by passing FCS through a 0.2 μm filter and mixing 1 ml of a 0.1 M disodium EDTA solution with 10 ml of filtered FCS.

Preparation of lung single cell suspensions

Following BAL, the pulmonary and systemic circulation is perfused with saline containing 5 mM EDTA to remove the intravascular pool of cells. One lung is used for histology, the other is used for the preparation of a cell suspension. The lung is thoroughly minced using iridectomy scissors and incubated for 30 min in digestion medium in a humidified incubator at 37°C and 5 % CO_2 . Organ fragments are resuspended, fresh digestion medium is added, and incubation is extended for another 15 min. After a final resuspension, very few organ debris are left. Samples are centrifuged and resuspended in calcium and magnesium-free PBS containing 10 mM EDTA at room temperature. Finally, the cells are subjected to RBC lysis, washed in FACS-EDTA, passed through a 50 μm cell strainer, and kept on ice until labeling. Cell counting is performed with a Z2 Beckman-Coulter particle counter (Beckman-Coulter).

Labeling of BAL-cells and lung single cell suspensions for flow cytometry

Cells are pre-incubated with Fc-receptor blocking antibody (anti-CD16/CD32, clone 2.4G2) to reduce non-specific binding. Monoclonal antibodies used to identify mouse DC populations are: biotinylated anti-CD11c (N418) and PE-conjugated anti-IA^b (AF6-120.1), followed by streptavidine-allophycocyanine (Sav-APC) (all from BD PharMingen). Isotype controls are PE-conjugated rat IgG_{2aK}, rat IgG_{2b} and armenian hamster IgG_{2K}. The following antibodies are used to stain mouse T-cell subpopulations: FITC-conjugated anti-CD4 (L3T4), FITC-conjugated anti-CD8 (Ly-2), and biotinylated anti-CD3 (145-2C11) monoclonal antibodies. The additional marker used for activation is anti-CD69 (H1.2F3). Biotinylated anti-CD3 is revealed by incubation with Sav-APC (all from BD Pharmingen). Monocytes/macrophages will be identified using FSC/SSC and CD11c staining.

As a last step before analysis, cells are incubated with 7-amino-actinomycin (7-AAD or viaprobe, BD Pharmingen) for dead cell exclusion. All labeling reactions are performed on ice in FACS-EDTA buffer.

Flow cytometry data acquisition is performed on a dual-laser FACS VantageTM flow cytometer running CELLQuestTM software (Becton Dickinson). FlowJo software (www.Treestar.com) is used for data analysis.

Histology

After clamping the main bronchus of the lung excised for the preparation of a cell suspension, fixative (4% paraformaldehyde in PBS) is gently infused through the tracheal cannula by a continuous-release pump under pressure and volume controlled conditions (12 ml/hour, 3 psi, 10 min/lung). The lung is resected and fixed for an additional 4 h. After routine paraffin embedding, 3 µm sections are stained with hematoxylin and eosin (H&E) (Klinipath) and examined by light microscopy for histological changes.

Quantification of emphysema is performed in a blinded fashion using a Zeiss KS400 Image Analyzer system running a custom-made morphometry program. The mean linear intercept (L_m) is measured for each mouse from 10 random fields by means of a 100 x 100 µm grid [15]. The mean alveolar surface (A_m) is also measured in 10 random fields per mouse. The number of alveolar wall breaks is counted per field and divided by the number of alveolar spaces in that field. The mean number of breaks/alveolar space is calculated from 10 random fields per mouse.

RESULTS:**1. Effect of anti-IFN-gamma mAb treatment on circulating TNF-alfa levels induced in smoking animals**

Proinflammatory cytokines, particularly IL-1beta and TNF, may amplify the inflammatory response in COPD and be linked to disease severity. TNF-alfa is present in high concentrations in the sputum of COPD patients. Serum concentration of TNF-alfa are increased in weight-losing COPD patients, suggesting that it may play a role in the cachexia of severe COPD. TNF-alfa inhibits the expression of skeletal muscle proteins via activation of NF-kB. This suggests that inhibitors of TNF-alfa might be useful in reversing the skeletal wasting seen in COPD as well as reducing the airway inflammatory response (Barnes, 2003; Oudijk, 2003)

In our experiment, inflammation in the smoking animals is evidenced by increased levels of TNF-alfa in the circulation of the smoking mice, as compared to the non-smoking animals. TNF alfa is detectable in the smoking animals after 3 to 4 weeks of smoking. TNF-alfa levels are decreased by treatment of the animals with anti-IFNgamma mAb, as shown in figure 1.

Our results demonstrate that blocking IFNgamma has an effect on the systemic levels of TNF-alfa. Thus, these data demonstrate that blocking IFN-gamma has an effect on the inflammatory response induced by smoking in these animals.

REFERENCES

- Aaron SD, Angel JB, Lunau M, et al. 2001. Granulocyte inflammatory markers and airway infection during acute exacerbation of chronic obstructive pulmonary disease. *Am. J. Resp. Crit. Care Med.* 163:349-55
- Alexander J., Sidney J., Southwood S., Ruppert J., Oseroff C., Maewal A., Snoke K., Serra HM., Kubo RT., Sette A., et al. (1994). Development of high potency universal DR-restricted helper epitopes by modification of high affinity DR-blocking peptides. *Immunity*, 1(9): 751-761.
- Ausubel et al. (1987) (1992) (1993) *Current Protocols in Molecular Biology*. Wiley Interscience, N.Y.
- Balbi B., Di Stefano A., Capelli A., Balbo P., Lusuardi M., Ioli F. and Donner C.F. T1 and T2 cytokines in bronchoalveolar lavage of patients with COPD and with asthma. International Conference of the American Thoracic Society, 2002.
- Barr I.G. and Mitchell G.F. (1996). ISCOMs (immunostimulating complexes): the first decade. *Immunol. and Cell Biol.* 74: 8-25.
- Barnes P.J. (2000). Chronic Obstructive Pulmonary Disease. *NEJM*, 343: 269-80.
- Barnes P.J. (2001). New treatments for chronic obstructive pulmonary disease. *Curr Opin Pharmacology*, 1: 217-222.
- Barnes P.J. (2002). New treatments for COPD. *Nature reviews Drug discovery*, 1: 437- 446.
- Barnes PJ (2003). New concepts in chronic obstructive pulmonary disease. *Annu. Rev. Med.*, 54:113-29.
- Baughman R.P. (2002). Therapeutic options for sarcoidosis: new and old. *Curr Opin Pulm Med*, 8: 464-469.

Billiau A. (1996) Interferon- γ : biology and role in pathogenesis. *Advances in Immunology* 62: 61-130.

Campbell A.M. (1984) *Monoclonal Antibody Technology Techniques in Biochemistry and Molecular Biology*, Elsevier Science Publishers, Amsterdam, The Netherlands.

Chackerian B., Lenz P., Lowy D.R. and Schiller J.T. (2002). Determinants of autoantibody induction by conjugated Papillomavirus virus-like particles. *J Immunol*, 169: 6120-6126.

Ciapponi L., Maione D., Scoumanne A., Costa P., Hansen M.B., Svenson M., Bendtzen K., Alonzi T., Paonessa G., Cortese R., Ciliberto G. and Savino R. (1997). Induction of interleukin-6 (IL-6) autoantibodies through vaccination with an engineered IL-6 receptor antagonist. *Nature Biotechnol*, 15: 997-1001.

Colligan J.E., Kruisbeek A.M., D.H. Margulies D.H. (1992) (1993) *Current Protocols in Immunology*, Green Publishing Association and Wiley Interscience, N.Y.

Contreras C.E., Ploton I.N., Siliciano R.F., Karp C.L., Viscidi R. and Kumar N. (1998). Mapping of specific and promiscuous HLA-DR-restricted T-cell epitopes on the *Plasmodium falciparum* 27-kilodalton sexual stage-specific antigen. *Infect Immun*, 66(8): 3579-90.

Casolaro MA, Bernaudin JF, Saltini C, Ferrans VJ, Crystal RG. (1988). Accumulation of Langerhans' cells on the epithelial surface of the lower respiratory tract in normal subjects in association with cigarette smoking. *Am Rev Respir Dis*, 137:406-11.

Dahm I., Jensen M.R., Hindersson P., Elsnér H.I., Mouritsen S. (1996). Breaking of B cell tolerance toward a highly conserved self protein. *J Immunol* 157(11): 4796-804.

Dalum I., Butler D.M., Jensen M.R., Hindersson P., Steinaa L., Waterston A.M., Grell S.N., Feldmann M., Elsner H.I. and Mouritsen S. (1999). Therapeutic antibodies elicited by immunization against TNF- α . *Nature Biotechnol*, 17: 666-669.

de Boer W.I. (2002). Cytokines and therapy in COPD. *CHEST*, 121: 209S-218S.

Donnelly J.J., Ulmer J.B., and Liu M.A. (1997a). DNA vaccines. *Life Sci.*, 60(3): 163-172.

Donnelly J.J., Ulmer J.B., Shiver J.W. and Liu M.A. (1997). DNA vaccines. *Annu. Rev. Immunol.*, 15: 617-648.

Doolan D.L., Southwood S., Chesnut R., Appella E., Gomez E., Richards A., Higashimoto Y.I., Maewal A., Sidney J., Gramzinski R.A., Mason C., Koech D., Hoffman S.L. and Sette A. (2000). HLA-DR-promiscuous T cell epitopes from *Plasmodium falciparum* pre-erythrocytic-stage antigens restricted by multiple HLA alleles. *J Immunol.*, 165: 1123-1137.

Dustin M.L., Singer K.H., Tuck D.T. and T.A. Springer (1988) Adhesion of T lymphoblasts to epidermal keratinocytes is regulated by interferon- γ and is mediated by intercellular adhesion molecule 1 (ICAM-1). *J. Exp. Med.* 167 : 1323-1340.

Fernando G.J., Tindle R.W. and Frazer I.H. (1995). T-helper epitopes of the E7 transforming protein of cervical cancer associated human papillomavirus type 18 (HPV18). *Virus Res*, 36(1): 1-13.

Freedman A.S., Freeman G.J., Rhyhart K. and L.M. Nadler (1991) Selective induction of B7/BB-1 on interferon-gamma stimulated monocytes: a potential mechanism for amplification of T cell activation through the CD28 pathway. *Cell. Immunol.* 137 : 429-437.

Friedl-Hajek R., Spangfort M.D., Schou C., Breiteneder H., Yssel H. and Joost van Neerven R.J. (1999). Identification of a highly promiscuous and an HLA-allele specific T-cell epitope in the birch major allergen Bet v 1: HLA restriction, epitope mapping and TCR sequence comparisons. *Clin Exp Allergy*, 29(4): 478-87.

Gaudebout P., Zeliszewski D., Golvano J.J., Pignal C., Le Gac S., Borrás-Cuesta F. and Sterkers G. (1997). Binding analysis of 95 HIV gp120 peptides to HLA-DR1101 and -DR0401 evidenced many HLA-class II binding regions on gp120 and suggested several promiscuous regions. *J Acquir Immune Defic Syndr Hum Retrovirol*, 14(2): 91-101.

Gish W. and States D.J. (1993). Identification of protein coding regions by database similarity search. *Nature Genet.*, 3:266-272.

Gonzalez S., Alvarez A., Caballero E., Vina L., Guillen G. and Silva R. (2000). P64k meningococcal protein as immunological carrier for weak immunogens. *Scand J Immunol*, 52(2): 113-116.

Harlow E., David L. et al. (1988) *Antibodies: A Laboratory Manual*. Cold Spring Harbor Publications.

Hautamaki R.D., Kobayashi D.K., Senior R.M. and Shapiro S.D. (1997). Requirement for macrophage elastase for cigarette smoke-induced emphysema in mice. *Science*, 277(5334):2002-4.

Holliger P., T. Prospero, and G. Winter (1993) Diabodies: small bivalent and bispecific antibody fragments. *Proc. Natl. Acad. Sci. USA* 90(14):6444.

Hurle M.R. and M. Gross (1994) Protein engineering techniques for antibody humanization. *Curr. Opin. Biotech.* 5:428-433.

Iliades P., Kortt A.A., and P. Hudson (1997) Triabodies: single chain Fv fragments without linker form trivalent trimers. *FEBS Letters* 409:437-441.

Kaplan N.M., Palmer B.F. and Weissler J.C. (2000). Syndromes of Severe Asthma. *Am Med Sci*, 319(3): 166-176.

Karlsson R., Michaelsson A. and Mattsson L. (1991). Kinetic analysis of monoclonal antibody-antigen interactions with a new biosensor based analytical system. *J Immunol Methods*, 145: 129-240.

Keatings VM, Collins PD, Scott DM, Barnes PJ. 1996. Differences in interleukin-8 and tumor necrosis factor- α in induced sputum from patients with chronic obstructive pulmonary disease or asthma. *Am. J. Resp. Crit. Care Med.* 153:530-34

Kilgus J., Jardetzky T., Gorga J.C., Trzeciak A., Gillesen D. and Sinigaglia F. (1991). Analysis of the permissive association of a malaria T cell epitope with DR molecules. *J Immunol*, 146(1): 307-15.

Kim S.K., Ragupathi G., Capello S., Kagan E. and Livingston P.O. (2001). Effect of immunological adjuvant combinations on the antibody and T cell response to vaccination with MUC1-KLH and GD3-KLH conjugates. *Vaccine*, 19: 530-537.

Kim Y-M. and Son K. (1996). A nitric oxide production assay for interferon- γ . *J Immunol Methods*, 198: 203-209.

Kobayashi H., Wood M., Song Y., Appella E. and Celis E. (2000). Defining promiscuous MHC class II helper T-cell epitopes for the HER2/neu tumor antigen. *Cancer Res*, 60(18): 5228-36.

Kortt A., Lah M., Oddie G., Gruen C., Burns J., Pearce L., Atwell J., McCoy A., Howlet G., Metzger D., Webster R. and P. Hudson (1997). Single-chain Fv fragments of anti-neuramidase antibody NC10 containing five- and ten-residue linkers form dimers and with zero-residue linker a trimer. *Prot. Eng.* 10:423-.

Köhler G. and C. Milstein (1975) Continuous cultures of fused cells secreting antibody of predefined specificity. *Nature* 256:495.

Launois P., Deleys R., Niang M.N., Drowart A., Andrien M., Dierckx P., Cartel J.L., Sarthou J.L., Van Vooren J.P. and Hyügen K. (1994). T-cell-epitope mapping of the

major secreted mycobacterial antigen Ag85A in tuberculosis and leprosy. *Infect Immun*, 62(9): 3679-87.

Leckie M.J., Bryan S.A., Hansel T.T. and Barnes P.J. (2000). Novel therapy for COPD. *Exp Opin Invest Drugs*, 9: 3-23.

Lethbridge M.W.G, Allenby M.I., Jensen M.W., Kemeny D.M. and O'Connor B.J. (2002). Airway T lymphocytes expressing IFN- γ are reduced in smokers with chronic obstructive pulmonary disease (COPD). Annual Congress 2002 of the European Respiratory Society- Stockholm.

Lewis J.A. (1995). A sensitive bioassay for interferons. *J Immunol Methods*, 185: 9-17.

Li Y., Wang M-N., Li H., King K.D., Bassi R., Sun H., Santiago A., Hooper A.T., Bohlen P. and Hicklin D.J. (2002). Active immunization against the vascular endothelial growth factor receptor flk1 inhibits tumor angiogenesis and metastasis. *J Exp Med*, 195(12): 1575-1584.

Lorsbach R.B., Murphy W.J., Lowenstein C.J., Snyder S.H. and S.W. Russel (1993) Expression of the nitric oxide synthase gene in mouse macrophages activated for tumor cell killing. *J. Biol. Chem.* 268 : 1908-1913.

Lowenadler B. and Lycke N. (1994). Fusion proteins with heterologous T helper epitopes. Recombinant *E. coli* heat-stable enterotoxin proteins. *Int Rev Immunol*, 11(2): 103-11.

Majori M., Corradi M., Caminati A., Giancarlo C., Bertacco S. and Pesci A. (1999). Predominant TH1 cytokine pattern in peripheral blood from patients with COPD. *J Allergy Clin Immunol*, 103: 458-62.

Mannino D.M. (2002). COPD: Epidemiology, prevalence, morbidity and mortality, and disease heterogeneity. *CHEST*, 121: 121S-126S.

Malmqvist M. (1999). BIACORE: an affinity biosensor system for characterization of biomolecular interactions. *Biochemical Society Transactions*, Vol 27.

Meager A. (2002). Biological assays for interferons. *J Immunol Methods*, 261 (1-2): 21-36.

Mendez MJ., Green L.L., Corvalan J.R., Jia X.C., Maynard-Currie C.E., Yang X.D., Gallo M.L., Louie D.M., Lee D.V., Erickson K.L., Luna J., Roy C.M., Abderrahim H., Kirschenbaum F., Noguchi M., Smith D.H., Fukushima A., Hales J.F., Klapholz S., Finer M.H., Davis C.G., Zsebo K.M. and Jakobovits A. (1997). Functional transplant of megabase human immunoglobulin loci recapitulates human antibody response in mice. *Nat Genet*, 15(2):146-56.

Möllers M., Aries S.P., Drömann D., Mascher B., Braun J. and Dalhoff K. (2001). Intracellular cytokine repertoire in different T cell subsets from patients with sarcoidosis. *Thorax*, 56: 487-493.

Morein B. et al. (1995). *Clin. Immunother.* 3: 461-475.

Muller R. (1983) Determination of affinity and specificity of anti-hapten antibodies by competitive radioimmunoassay. *Methods Enzymol.* 92: 589-601.

Mustafa A.S., Shaban F.A., Abal A.T., Al-attiyah R., Wiker H.G., Lundin K.E., Oftung F. and Huygen K. (2000). Identification and HLA restriction of naturally derived Th1-cell epitopes from the secreted Mycobacterium tuberculosis antigen 85B recognized by antigen-specific human CD4+ T-cell lines. *Infect. Immun.*, July: 3933-3940.

NIH, National heart, lung and blood institute (1995). Sarcoidosis. NIH publication No. 95-3093, reprinted July 1995.

NIH, National heart, lung and blood institute (1997). The Lungs in health and disease. NIH Publication No. 97-3279, August 1997.

NHLBI/WHO. Executive Summary, Global Strategy for the diagnosis, management and prevention of chronic obstructive pulmonary disease. Workshop Report.
Pace J.L., Russell S.W., Torres B.A., Johnson H.M. and P.W. Gray (1983) Recombinant mouse γ interferon induces the priming step in macrophage activation for tumor cell killing. *J.Immunol.* 130 : 2011-2013.

Oudijk E-JD, Lammers J-W J, Koenderman L. (2003) Systemic inflammation in chronic obstructive pulmonary disease. *Eur Respir J*, 22(S46): 5s-13s

Papiris S., Kotanidou A., Malagari K. and Roussos C. (2002). Clinical review: severe asthma. *Crit Care*, 6(1): 30-44.

Pauwels RA, Buist AS, Calverley PM, Jenkins CR, Hurd SS.(2001) Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. NHLBI/WHO Global Initiative for Chronic Obstructive Lung Disease (GOLD) Workshop summary. *Am J Respir Crit Care Med*, 163:1256-76.

Retamales I, Elliott WM, Meshi B, Coxson HO, Pare PD, Sciruba FC, et al. (2001). Amplification of inflammation in emphysema and its association with latent adenoviral infection. *Amer J Respir Crit Care Med*, 164: 469-473.

Poljak R.J. (1994) Production and structure of diabodies. *Structure* 2:1121-1123.

Richard M., Grecis R.K., Humphreys N.E., Renauld J-C. and Van Snick J. (2000). Anti-IL-9 vaccination prevents worm expulsion and blood eosinophilia in *Trichuris muris*-infected mice. *PNAS*, 97(2): 767-772.

Roguska M.A., Pedersen J.T., Keddy C.A., Henry A.H., Searle S.J., Lambert J.M., Goldmacher V.S., Blättler W.A., Rees A.R. and B.C. Guild (1994) Humanization of

murine monoclonal antibodies through variable domain resurfacing. *Proc. Natl. Acad. Sci. USA* 91 : 969-973.

Rossmann M.D. (2001). Chronic Beryllium Disease: A hypersensitivity disorder. *Appl Occup Environ Hygiene*, 16(5): 615-618.

Sad S., Rao K., Arora R., Talwar G.P. and Raghupathy R. (1992). Bypass of carrier-induced epitope-specific suppression using a T-helper epitope. *Immunology*, 76: 599-603.

Sangiulio F., D'Apice M.R., Bruscia E., Lucidi V. and Novelli G. (2002). Towards the pharmacogenomics of cystic fibrosis. *Pharmacogenomics*, 3(1): 75-87.

Salini C. (2001). Beryllium Disease. *Am J Med Sci*, 321(1): 89-98.

Shigehara K., Shijubo N., Ohmichi M., Takahashi R., Kon S., Okamura H., Kurimoto M., Hiraga Y., Tatsuno T., Abe S. and Sato N. (2001). IL-12 and IL-18 are increased and stimulate IFN- γ production in sarcoid lungs. *J Immunol*, 166: 642-649.

Snapper C.M. and W.E. Paul (1987) Interferon- γ and B cell stimulatory factor-1 reciprocally regulate Ig isotype production. *Science* 236 : 944-947.

Steinman R.M., Noguiera N., Witmer M.D., Tydings J.G. and I.S. Mellman (1980) Lymphokine enhances the expression and synthesis of Ia antigens on cultured mouse peritoneal macrophages. *J. exp. Med.* 152 : 1248-1261.

Stirling R.G. and Chung K.F. (2001). Severe Asthma: Definition and mechanisms. *Allergy*, 56: 825-840.

Storni T., Lechner F., Erdmann I., Bächli T., Jegerlehner A., Dumrese T., Kündig T.M., Ruedi C. and Bachmann M.F. (2002). Critical role for activation of antigen-presenting cells in priming of cytotoxic T cell responses after vaccination with virus-like particles. *J Immunol*, 168: 2880-2886.

Talwar G.P. (1999). Vaccines and passive immunological approaches for the control of fertility and hormone-dependent cancers. *Immunological Reviews*, 171: 173-192.

Udono H. and Srivastava P.K. (1993). Heat shock protein 70-associated peptides elicit specific cancer immunity. *J Exp Med*, 178(4): 1391-6.

Vernersson M., Ledin A., Johansson J. and Hellman L. (2002). Generation of therapeutic antibody responses against IgE through vaccination. *FASEB J*, 16: 875-877.

Wang Z., Zheng T., Zhu Z., Homer R.J., Riese R., Chapman H.A., Shapiro S.D. and Elias J.A. (2000). Interferon γ induction of pulmonary emphysema in the adult murine lung. *J Exp Med*, 192(11): 1587-1599.

Wenzel S.E., Szefer S.J., Leung D.Y., Sloan S.I., Rex M.D. and Martin R.J. (1997). Bronchoscopic evaluation of severe asthma. Persistent inflammation associated with high dose glucocorticoids. *Am J Respir Crit Care Med*, 156(3 Pt 1):737-43.

Zhu Z., Zapata G., Shalaby R., Snedecor B., Chen H. and P. Carter (1996) High level secretion of a humanized bispecific diabody from *Escherichia coli*. *Biotechnology* 14:192-196.

CLAIMS

1. Use of an IFN γ neutralizing molecule for the manufacture of a medicament for preventing or treating a T1 inflammatory lung disease.
2. Use according to claim 1 wherein said T1 inflammatory lung disease is selected from the group consisting of COPD, emphysema, chronic bronchitis, bronchiolitis, severe asthma, sarcoidosis, berylliosis and cystic fibrosis.
3. Use according to claims 1-2, wherein said molecule is an anti-IFN γ antibody or a fragment thereof.
4. Use according to claim 3, wherein said antibody is a monoclonal antibody.
5. Use according to claim 4, wherein said antibody is the antibody D9D10 or a fragment thereof.
6. Use according to claim 3, wherein said antibody is a humanized monoclonal antibody.
7. Use according to claim 6, wherein said antibody is a humanized D9D10 antibody.
8. Use of immunogenic IFN γ for the manufacture of a medicament for preventing or treating a T1 inflammatory lung disease.
9. Use according to claim 8 wherein said T1 inflammatory lung disease is selected from the group consisting of COPD, emphysema, chronic bronchitis, bronchiolitis, severe asthma, sarcoidosis, berylliosis and cystic fibrosis.
10. Use according to claim 8, wherein immunogenic IFN γ is a modified IFN γ .
11. Use according to claim 10, wherein the IFN γ protein is modified to immunogenic IFN γ by crosslinking IFN γ to an immunogenic non-self carrier protein.

12. Use according to claim 10, wherein the IFN γ protein is modified to immunogenic IFN γ by introducing at least one foreign T-cell epitope by insertion and/or substitution and/or addition and/or conjugation.
13. Use according to claim 8, wherein immunogenic IFN γ is a non-self IFN γ administered with or without adjuvant.
14. Use according to claim 8, wherein immunogenic IFN γ is autologous IFN γ in combination with an adjuvant.
15. Use of at least one autologous antigen presenting cell loaded with IFN γ for the manufacture of a medicament for preventing or treating a T1 inflammatory lung disease.
16. Use according to claim 16, wherein the antigen presenting cell is a dendritic cell.

10/550175

1/1





Figure 1

Original (for SUBMISSION) - printed on Friday, 26 March, 2004 01:46:23 PM

0-1	Form - PCT/RO/134 (EASY) Indications Relating to Deposited Microorganism(s) or Other Biological Material (PCT Rule 13bls)	
0-1-1	Prepared using	epoline® online filing PCT plug-in (updated 26.03.2004)
0-2	International Application No.	PCT/EP2004 / 050375
0-3	Applicant's or agent's file reference	149 PCT

1	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
1-1	page	15
1-2	line	30
1-3	Identification of Deposit	
1-3-1	Name of depositary institution	DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH
1-3-2	Address of depositary institution	Mascheroder Weg 1b, D-38124 Braunschweig, Germany
1-3-3	Date of deposit	28 August 2001 (28.08.2001)
1-3-4	Accession Number	DSMZ ACC2521
1-4	Additional Indications	NONE
1-5	Designated States for Which Indications are Made	all designated States
1-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE

FOR RECEIVING OFFICE USE ONLY

0-4	This form was received with the international application: (yes or no)	YES
0-4-1	Authorized officer	

FOR INTERNATIONAL BUREAU USE ONLY

0-5	This form was received by the international Bureau on:	
0-5-1	Authorized officer	

INTERNATIONAL SEARCH REPORT

tional Application No
EP2004/050375

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07K16/24 A61K39/395 A61K39/385 A61K39/00 A61P37/02 A61P11/00 A61P37/08		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07K A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, BIOSIS, MEDLINE, PAJ, WPI Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DENIS M ET AL: "MURINE HYPERSENSITIVITY PNEUMONITIS BIDIRECTIONAL ROLE OF INTERFERON-GAMMA" CLINICAL AND EXPERIMENTAL ALLERGY, vol. 22, no. 8, 1992, pages 783-792, XP009018515 ISSN: 0954-7894	1-4,8,9
Y	abstract page 790, left-hand column, paragraph 2 page 784, left-hand column page 789, left-hand column, paragraph 2	1-16
X	WO 01/34180 A (ZIESCHE ROLF ;BLOCK LUTZ HENNING (DE)) 17 May 2001 (2001-05-17) page 2; claims 1,2 ----- -/-	8-10
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *&* document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
22 July 2004		02/09/2004
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Lechner, O

INTERNATIONAL SEARCH REPORT

onal Application No

EP2004/050375

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 02/069996 A (LOVELACE RESPIRATORY RES INST ;TESFAIGZI YOHANNES (US)) 12 September 2002 (2002-09-12) abstract page 1, line 26 - page 2, line 13 page 4, lines 23-26 page 5, lines 1-11 page 7, lines 4-6 page 11, lines 22-30</p>	8-10
Y	<p>WANG ZHONGDE ET AL: "Interferon gamma induction of pulmonary emphysema in the adult murine lung." JOURNAL OF EXPERIMENTAL MEDICINE, vol. 192, no. 11, 4 December 2000 (2000-12-04), pages 1587-1599, XP002257504 ISSN: 0022-1007 the whole document</p>	1-16
Y	<p>ROBINSON B W ET AL: "Gamma interferon is spontaneously released by alveolar macrophages and lung T lymphocytes in patients with pulmonary sarcoidosis." THE JOURNAL OF CLINICAL INVESTIGATION. UNITED STATES MAY 1985, vol. 75, no. 5, May 1985 (1985-05), pages 1488-1495, XP009018516 ISSN: 0021-9738 the whole document</p>	1-16
Y	<p>FROYEN G ET AL: "Potential therapeutic use of antibodies directed towards HuIFN-gamma" BIOTHERAPY, KLUWER ACADEMIC PUBLISHERS, DORDRECHT, NL, vol. 10, 1997, pages 49-57, XP002067597 ISSN: 0921-299X the whole document</p>	1-16
Y	<p>WO 95/05849 A (ELSNER HENRIK ;MOURITSEN & ELSNER A S (DK); MOURITSEN SOEREN (DK)) 2 March 1995 (1995-03-02) cited in the application the whole document</p>	1-16

-/--

INTERNATIONAL SEARCH REPORT

International Application No
/EP2004/050375

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	DALUM I ET AL: "Therapeutic antibodies elicited by immunization against TNF-alpha." NATURE BIOTECHNOLOGY. UNITED STATES JUL 1999, vol. 17, no. 7, July 1999 (1999-07), pages 666-669, XP000910155 ISSN: 1087-0156 cited in the application the whole document	1-16
A	CARUSO A ET AL: "Natural antibodies to IFN-gamma in man and their increase during viral infection." JOURNAL OF IMMUNOLOGY (BALTIMORE, MD.: 1950) UNITED STATES 15 JAN 1990, vol. 144, no. 2, 15 January 1990 (1990-01-15), pages 685-690, XP002257507 ISSN: 0022-1767 the whole document	1-16
A	BEERS-MH AND BERKOW-R (EDS): "The merck manual of diagnosis and therapy" 1999, MERCK RESEARCH LABORATORIES , XP002257638 page 628, right-hand column, last paragraph	1-16

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

/EP2004/050375

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0134180	A	17-05-2001	AU 1392301 A	06-06-2001
			CA 2387089 A1	17-05-2001
			WO 0134180 A2	17-05-2001
			EP 1227831 A2	07-08-2002
			JP 2003513933 T	15-04-2003
WO 02069996	A	12-09-2002	WO 02069996 A1	12-09-2002
WO 9505849	A	02-03-1995	AT 162723 T	15-02-1998
			AU 707083 B2	01-07-1999
			AU 7009198 A	30-07-1998
			AU 7608094 A	21-03-1995
			CA 2170236 A1	02-03-1995
			DE 69408342 D1	05-03-1998
			DE 69408342 T2	14-05-1998
			WO 9505849 A1	02-03-1995
			DK 752886 T3	04-05-1998
			EP 0752886 A1	15-01-1997
			ES 2112559 T3	01-04-1998
			GR 3026419 T3	30-06-1998
			JP 9505031 T	20-05-1997
			US 2002090379 A1	11-07-2002